



2/12




FIG.2

3/12




FIG.3

4/12




FIG.4

2017-04-12 14:00

5/12

201720" E44E400F




FIG.5

6/12




FIG.6

7/12




FIG.7

8/12




FIG.8

9/12

RAT JEJUNAL ABSORPTION OF ODV SUCCINATE AND INTERNAL MARKERS
[ODV SUCCINATE] = 50 ug/ml, PERFUSION BUFFER, pH6.8




FIG.9

INTESTINAL SITE-SPECIFIC ABSORPTION OF ODV SUCCINATE
[ODV SUCCINATE] = 50 ug/ml, PERFUSION BUFFER, pH6.8




FIG.10

10/12

RAT JEJUNAL ABSORPTION OF ODV FUMARATE AND INTERNAL MARKERS
[ODV FUMARATE] = 50 ug/ml, PERFUSION BUFFER, pH6.8




FIG.11

INTESTINAL SITE-SPECIFIC ABSORPTION OF ODV FUMARATE
[ODV FUMARATE] = 50 ug/ml, PERFUSION BUFFER, pH6.8




FIG.12

11/12

COMPARISON OF SITE-SPECIFIC ABSORPTION:
ODV FUMARATE vs ODV SUCCINATE




FIG.13

12/12

